


GRANISETRON HYDROCHLORIDE - granisetron hydrochloride tablet, film coated
COREPHARMA LLC.

DESCRIPTION

Granisetron hydrochloride tablets contain granisetron hydrochloride, an antiemetic agent. Chemically it is *endo*-N-(9-methyl-9-azabicyclo [3.3.1] non-3-yl)-1-methyl-1H-indazole-3-carboxamide hydrochloride with a molecular weight of 348.9 (312.4 free base). Its molecular formula is C₁₈H₂₄N₄O•HCl while its chemical structure is:



Granisetron hydrochloride is a white to off-white solid that is readily soluble in water and normal saline at 20°C.

Tablets for Oral Administration: Each white round film coated tablet contains 1.12 mg granisetron hydrochloride equivalent to granisetron, 1 mg. In addition, each tablet contains the following inactive ingredients: hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, sodium starch glycolate and titanium dioxide.

CLINICAL PHARMACOLOGY

Granisetron is a selective 5-hydroxytryptamine₃ (5-HT₃) receptor antagonist with little or no affinity for other serotonin receptors, including 5-HT₁; 5-HT_{1A}; 5-HT_{1B/C}; 5-HT₂; for alpha₁-, alpha₂-, or beta-adrenoreceptors; for dopamine-D₂; or for histamine-H₁; benzodiazepine; picrotoxin or opioid receptors.

Serotonin receptors of the 5-HT₃ type are located peripherally on vagal nerve terminals and centrally in the chemoreceptor trigger zone of the area postrema. During chemotherapy that induces vomiting, mucosal enterochromaffin cells release serotonin, which stimulates 5-HT₃ receptors. This evokes vagal afferent discharge, inducing vomiting. Animal studies demonstrate that, in binding to 5-HT₃ receptors, granisetron blocks serotonin stimulation and subsequent vomiting after emetogenic stimuli such as cisplatin. In the ferret animal model, a single granisetron injection prevented vomiting due to high-dose cisplatin or arrested vomiting within 5 to 30 seconds.

In most human studies, granisetron has had little effect on blood pressure, heart rate or ECG. No evidence of an effect on plasma prolactin or aldosterone concentrations has been found in other studies.

Following single and multiple oral doses, granisetron hydrochloride tablets slowed colonic transit in normal volunteers. However, granisetron hydrochloride had no effect on oro-cecal transit time in normal volunteers when given as a single intravenous (IV) infusion of 50 mcg/kg or 200 mcg/kg.

Pharmacokinetics

In healthy volunteers and adult cancer patients undergoing chemotherapy, administration of granisetron hydrochloride tablets produced mean pharmacokinetic data shown in **Table 1**.

Table 1 Pharmacokinetic Parameters (Median [range]) Following Granisetron Hydrochloride Tablets

	Peak Plasma Concentration (ng/mL)	Terminal Phase Plasma Half-Life (h)	Volume of Distribution (L/kg)	Total Clearance (L/h/kg)
Cancer Patients 1 mg bid, 7 days (n=27)	5.99 [0.63 to 30.9]	N.D.*	N.D.	0.52 [0.09 to 7.37]
Volunteers Single 1 mg dose (n=39)	3.63 [0.27 to 9.14]	6.23 [0.96 to 19.9]	3.94 [1.89 to 39.4]	0.41 [0.11 to 24.6]

*Not determined after oral administration; following a single intravenous dose of 40 mcg/kg, terminal phase half-life was determined to be 8.95 hours.

N.D. Not determined.

Absorption : When granisetron hydrochloride tablets were administered with food, AUC was decreased by 5% and C_{max} increased by 30% in non-fasted healthy volunteers who received a single dose of 10 mg.

Distribution : Plasma protein binding is approximately 65% and granisetron distributes freely between plasma and red blood cells.

Metabolism : Granisetron metabolism involves N-demethylation and aromatic ring oxidation followed by conjugation. *In vitro* liver microsomal studies show that granisetron's major route of metabolism is inhibited by ketoconazole, suggestive of metabolism mediated by the cytochrome P-450 3A subfamily. Animal studies suggest that some of the metabolites may also have 5-HT₃ receptor antagonist activity.

Elimination : Clearance is predominantly by hepatic metabolism. In normal volunteers, approximately 11% of the orally administered dose is eliminated unchanged in the urine in 48 hours. The remainder of the dose is excreted as metabolites, 48% in the urine and 38% in the feces.

Subpopulations

Gender : The effects of gender on the pharmacokinetics of granisetron hydrochloride tablets have not been studied. However, after intravenous infusion of granisetron hydrochloride, no difference in mean AUC was found between males and females, although males had a higher C_{max} generally.

In elderly and pediatric patients and in patients with renal failure or hepatic impairment, the pharmacokinetics of granisetron was determined following administration of intravenous granisetron hydrochloride.

Elderly : The ranges of the pharmacokinetic parameters in elderly volunteers (mean age 71 years), given a single 40 mcg/kg intravenous dose of granisetron hydrochloride injection, were generally similar to those in younger healthy volunteers; mean values were lower for clearance and longer for half-life in the elderly.

Renal Failure Patients : Total clearance of granisetron was not affected in patients with severe renal failure who received a single 40 mcg/kg intravenous dose of granisetron hydrochloride injection.

Hepatically Impaired Patients : A pharmacokinetic study with intravenous granisetron hydrochloride in patients with hepatic impairment due to neoplastic liver involvement showed that total clearance was approximately halved compared to patients without hepatic impairment. Given the wide variability in pharmacokinetic parameters noted in patients and the good tolerance of doses well above the recommended dose, dosage adjustment in patients with possible hepatic functional impairment is not necessary.

Pediatric Patients : A pharmacokinetic study in pediatric cancer patients (2 to 16 years of age), given a single 40 mcg/kg intravenous dose of granisetron hydrochloride injection, showed that volume of distribution and total clearance increased with age. No relationship with age was observed for peak plasma concentration or terminal phase plasma half-life. When volume of distribution and total clearance are adjusted for body weight, the pharmacokinetics of granisetron are similar in pediatric and adult cancer patients.

CLINICAL TRIALS

Chemotherapy-Induced Nausea and Vomiting

Granisetron hydrochloride tablets prevent nausea and vomiting associated with initial and repeat courses of emetogenic cancer therapy, as shown by 24-hour efficacy data from studies using both moderately-and highly-emetogenic chemotherapy.

Moderately Emetogenic Chemotherapy : The first trial compared granisetron hydrochloride tablets doses of 0.25 mg to 2 mg bid, in 930 cancer patients receiving, principally, cyclophosphamide, carboplatin, and cisplatin (20 mg/m² to 50 mg/m²). Efficacy was based on complete response (ie, no vomiting, no moderate or severe nausea, no rescue medication), no vomiting, and no nausea. **Table 2** summarizes the results of this study.

Table 2 Prevention of Nausea and Vomiting 24 Hours Post-Chemotherapy*

Efficacy Measures	Percentages of Patients Granisetron Hydrochloride Tablet Dose			
	0.25 mg bid (n=229) %	0.5 mg bid (n=235) %	1 mg bid (n=233) %	2 mg bid (n=233) %
Complete Response [†]	61	70 [‡]	81 ^{‡§}	72 [‡]
No Vomiting	66	77 [‡]	88 [‡]	79 [‡]
No Nausea	48	57	63 [‡]	54

*Chemotherapy included oral and injectable cyclophosphamide, carboplatin, cisplatin (20 mg/m² to 50 mg/m²), dacarbazine, doxorubicin, epirubicin.

†No Vomiting, no moderate or severe nausea, no rescue medication.

‡Statistically significant (P<0.01) vs. 0.25 mg bid.

§Statistically significant (P<0.01) vs. 0.5 mg bid.

Results from a second double-blind, randomized trial evaluating granisetron hydrochloride tablets 2 mg qd and granisetron hydrochloride tablets 1 mg bid were compared to prochlorperazine 10 mg bid derived from a historical control. At 24 hours, there was no statistically significant difference in efficacy between the two granisetron hydrochloride tablet regimens. Both regimens were statistically superior to the prochlorperazine control regimen (see **Table 3**).

Table 3 Prevention of Nausea and Vomiting 24 Hours Post-Chemotherapy *

Efficacy Measures	Percentages of Patients		
	Granisetron HCl Tablets 1 mg bid (n=354) %	Granisetron HCl Tablets 2 mg qd (n=343) %	Prochlorperazine [†] 10 mg bid (n=111) %
Complete Response [‡]	69 [§]	64 [§]	41
No Vomiting	82 [§]	77 [§]	48
No Nausea	51 [§]	53 [§]	35
Total Control [¶]	51 [§]	50 [§]	33

*Moderately emetogenic chemotherapeutic agents included cisplatin (20 mg/m² to 50 mg/m²), oral and intravenous cyclophosphamide, carboplatin, dacarbazine, doxorubicin.

†Historical control from a previous double-blind granisetron hydrochloride trial.

‡No vomiting, no moderate or severe nausea, no rescue medication.

§Statistically significant (P<0.05) vs. prochlorperazine historical control.

¶No vomiting, no nausea, no rescue medication.

Results from a granisetron hydrochloride tablets 2 mg qd alone treatment arm in a third double-blind, randomized trial, were compared to prochlorperazine (PCPZ), 10 mg bid, derived from a historical control. The 24-hour results for granisetron hydrochloride tablets 2 mg qd were statistically superior to PCPZ for all efficacy parameters: complete response (58%), no vomiting (79%), no nausea (51%), total control (49%). The PCPZ rates are shown in **Table 3**.

Cisplatin-Based Chemotherapy : The first double-blind trial compared granisetron hydrochloride tablets 1 mg bid, relative to placebo (historical control), in 119 cancer patients receiving high-dose cisplatin (mean dose 80 mg/m²). At 24 hours, granisetron

hydrochloride tablets 1 mg bid was significantly ($P < 0.001$) superior to placebo (historical control) in all efficacy parameters: complete response (52%), no vomiting (56%) and no nausea (45%). The placebo rates were 7%, 14%, and 7%, respectively, for the three efficacy parameters.

Results from a granisetron hydrochloride tablets 2 mg qd alone treatment arm in a second double-blind, randomized trial, were compared to both granisetron hydrochloride tablets 1 mg bid and placebo historical controls. The 24-hour results for granisetron hydrochloride tablets 2 mg qd were: complete response (44%), no vomiting (58%), no nausea (46%), total control (40%). The efficacy of granisetron hydrochloride tablets 2 mg qd was comparable to granisetron hydrochloride tablets 1 mg bid and statistically superior to placebo. The placebo rates were 7%, 14%, 7%, and 7%, respectively, for the four parameters.

No controlled study comparing granisetron injection with the oral formulation to prevent chemotherapy-induced nausea and vomiting has been performed.

Radiation-Induced Nausea and Vomiting

Total Body Irradiation : In a double-blind randomized study, 18 patients receiving granisetron hydrochloride tablets, 2 mg daily, experienced significantly greater antiemetic protection compared to patients in a historical negative control group who received conventional (non-5-HT₃ antagonist) antiemetics. Total body irradiation consisted of 11 fractions of 120 cGy administered over 4 days, with three fractions on each of the first 3 days, and two fractions on the fourth day. Granisetron hydrochloride tablets were given one hour before the first radiation fraction of each day.

Twenty-two percent (22%) of patients treated with granisetron hydrochloride tablets did not experience vomiting or receive rescue antiemetics over the entire 4-day dosing period, compared to 0% of patients in the historical negative control group ($P < 0.01$).

In addition, patients who received granisetron hydrochloride tablets also experienced significantly fewer emetic episodes during the first day of radiation and over the 4-day treatment period, compared to patients in the historical negative control group. The median time to the first emetic episode was 36 hours for patients who received granisetron hydrochloride tablets.

Fractionated Abdominal Radiation : The efficacy of granisetron hydrochloride tablets, 2 mg daily, was evaluated in a double-blind, placebo-controlled randomized trial of 260 patients. Granisetron hydrochloride tablets were given 1 hour before radiation, composed of up to 20 daily fractions of 180 to 300 cGy each. The exceptions were patients with seminoma or those receiving whole abdomen irradiation who initially received 150 cGy per fraction. Radiation was administered to the upper abdomen with a field size of at least 100 cm².

The proportion of patients without emesis and those without nausea for granisetron hydrochloride tablets, compared to placebo, was statistically significant ($P < 0.0001$) at 24 hours after radiation, irrespective of the radiation dose. Granisetron hydrochloride was superior to placebo in patients receiving up to 10 daily fractions of radiation, but was not superior to placebo in patients receiving 20 fractions.

Patients treated with granisetron hydrochloride tablets ($n=134$) had a significantly longer time to the first episode of vomiting (35 days vs. 9 days, $P < 0.001$) relative to those patients who received placebo ($n=126$), and a significantly longer time to the first episode of nausea (11 days vs. 1 day, $P < 0.001$). Granisetron hydrochloride provided significantly greater protection from nausea and vomiting than placebo.

INDICATIONS AND USAGE

Granisetron hydrochloride tablets are indicated for the prevention of:

- Nausea and vomiting associated with initial and repeat courses of emetogenic cancer therapy, including high-dose cisplatin.
- Nausea and vomiting associated with radiation, including total body irradiation and fractionated abdominal radiation.

CONTRAINDICATIONS

Granisetron hydrochloride tablets are contraindicated in patients with known hypersensitivity to the drug or any of its components.

PRECAUTIONS

Granisetron hydrochloride is not a drug that stimulates gastric or intestinal peristalsis. It should not be used instead of nasogastric suction. The use of granisetron hydrochloride in patients following abdominal surgery or in patients with chemotherapy-induced nausea and vomiting may mask a progressive ileus and/or gastric distention.

Drug Interactions

Granisetron does not induce or inhibit the cytochrome P-450 drug-metabolizing enzyme system *in vitro*. There have been no definitive drug-drug interaction studies to examine pharmacokinetic or pharmacodynamic interaction with other drugs; however, in humans, granisetron hydrochloride injection has been safely administered with drugs representing benzodiazepines, neuroleptics, and anti-

ulcer medications commonly prescribed with antiemetic treatments. Granisetron hydrochloride injection also does not appear to interact with emetogenic cancer chemotherapies. Because granisetron is metabolized by hepatic cytochrome P-450 drug-metabolizing enzymes, inducers or inhibitors of these enzymes may change the clearance and, hence, the half-life of granisetron. No specific interaction studies have been conducted in anesthetized patients. In addition, the activity of the cytochrome P-450 subfamily 3A4 (involved in the metabolism of some of the main narcotic analgesic agents) is not modified by granisetron hydrochloride *in vitro*. In *in vitro* human microsomal studies, ketoconazole inhibited ring oxidation of granisetron hydrochloride. However, the clinical significance of *in vivo* pharmacokinetic interactions with ketoconazole is not known. In a human pharmacokinetic study, hepatic enzyme induction with phenobarbital resulted in a 25% increase in total plasma clearance of intravenous granisetron hydrochloride. The clinical significance of this change is not known.

Carcinogenesis, Mutagenesis, Impairment of Fertility

In a 24-month carcinogenicity study, rats were treated orally with granisetron 1, 5 or 50 mg/kg/day (6, 30 or 300 mg/m²/day). The 50 mg/kg/day dose was reduced to 25 mg/kg/day (150 mg/m²/day) during week 59 due to toxicity. For a 50 kg person of average height (1.46 m² body surface area), these doses represent 4, 20, and 101 times the recommended clinical dose (1.48 mg/m², oral) on a body surface area basis. There was a statistically significant increase in the incidence of hepatocellular carcinomas and adenomas in males treated with 5 mg/kg/day (30 mg/m²/day, 20 times the recommended human dose based on body surface area) and above, and in females treated with 25 mg/kg/day (150 mg/m²/day, 101 times the recommended human dose based on body surface area). No increase in liver tumors was observed at a dose of 1 mg/kg/day (6 mg/m²/day, 4 times the recommended human dose based on body surface area) in males and 5 mg/kg/day (30 mg/m²/day, 20 times the recommended human dose based on body surface area) in females. In a 12-month oral toxicity study, treatment with granisetron 100 mg/kg/day (600 mg/m²/day, 405 times the recommended human dose based on body surface area) produced hepatocellular adenomas in male and female rats while no such tumors were found in the control rats. A 24-month mouse carcinogenicity study of granisetron did not show a statistically significant increase in tumor incidence, but the study was not conclusive.

Because of the tumor findings in rat studies, granisetron hydrochloride should be prescribed only at the dose and for the indication recommended (see **INDICATIONS AND USAGE** and **DOSAGE AND ADMINISTRATION**).

Granisetron was not mutagenic in *in vitro* Ames test and mouse lymphoma cell forward mutation assay, and *in vivo* mouse micronucleus test and *in vitro* and *ex vivo* rat hepatocyte UDS assays. It, however, produced a significant increase in UDS in HeLa cells *in vitro* and a significant increased incidence of cells with polyploidy in an *in vitro* human lymphocyte chromosomal aberration test.

Granisetron at oral doses up to 100 mg/kg/day (600 mg/m²/day, 405 times the recommended human dose based on body surface area) was found to have no effect on fertility and reproductive performance of male and female rats.

Pregnancy

Teratogenic Effects : Pregnancy Category B. Reproduction studies have been performed in pregnant rats at oral doses up to 125 mg/kg/day (750 mg/m²/day, 507 times the recommended human dose based on body surface area) and pregnant rabbits at oral doses up to 32 mg/kg/day (378 mg/m²/day, 255 times the recommended human dose based on body surface area) and have revealed no evidence of impaired fertility or harm to the fetus due to granisetron. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Nursing Mothers : It is not known whether granisetron is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when granisetron hydrochloride is administered to a nursing woman.

Pediatric Use : Safety and effectiveness in pediatric patients have not been established.

Geriatric Use : During clinical trials, 325 patients 65 years of age or older received granisetron hydrochloride tablets; 298 were 65 to 74 years of age, and 27 were 75 years of age or older. Efficacy and safety were maintained with increasing age.

ADVERSE REACTIONS

Chemotherapy-Induced Nausea and Vomiting

Over 3700 patients have received granisetron hydrochloride tablets in clinical trials with emetogenic cancer therapies consisting primarily of cyclophosphamide or cisplatin regimens.

In patients receiving granisetron hydrochloride tablets 1 mg bid for 1, 7 or 14 days, or 2 mg qd for 1 day, adverse experiences reported in more than 5% of the patients with comparator and placebo incidences are listed in **Table 4**.

Table 4 Principal Adverse Events in Clinical Trials

	Percent of Patients With Event
--	--------------------------------

	Granisetron HCl* Tablets 1 mg bid (n=978)	Granisetron HCl* Tablets 2 mg qd (n=1450)	Comparator[†] (n=599)	Placebo (n=185)
Headache [‡]	21%	20%	13%	12%
Constipation	18%	14%	16%	8%
Asthenia	14%	18%	10%	4%
Diarrhea	8%	9%	10%	4%
Abdominal pain	6%	4%	6%	3%
Dyspepsia	4%	6%	5%	4%

*Adverse events were recorded for 7 days when granisetron hydrochloride tablets were given on a single day and for up to 28 days when granisetron hydrochloride tablets were administered for 7 to 14 days.

[†]Metoclopramide/
dexamethasone;
phenothiazines/
dexamethasone;
dexamethasone alone;
prochlorperazine

[‡]Usually mild to moderate in severity.

Other adverse events reported in clinical trials were:

Gastrointestinal : In single-day dosing studies in which adverse events were collected for 7 days, nausea (20%) and vomiting (12%) were recorded as adverse events after the 24-hour efficacy assessment period.

Hepatic : In comparative trials, elevation of AST and ALT (>2 times the upper limit of normal) following the administration of granisetron hydrochloride tablets occurred in 5% and 6% of patients, respectively. These frequencies were not significantly different from those seen with comparators (AST: 2%; ALT: 9%).

Cardiovascular : Hypertension (1%); hypotension, angina pectoris, atrial fibrillation, and syncope have been observed rarely.

Central Nervous System : Dizziness (5%), insomnia (5%), anxiety (2%), somnolence (1%). One case compatible with, but not diagnostic of, extrapyramidal symptoms has been reported in a patient treated with granisetron hydrochloride tablets.

Hypersensitivity : Rare cases of hypersensitivity reactions, sometimes severe (eg, anaphylaxis, shortness of breath, hypotension, urticaria) have been reported.

Other : Fever (5%). Events often associated with chemotherapy also have been reported: leukopenia (9%), decreased appetite (6%), anemia (4%), alopecia (3%), thrombocytopenia (2%).

Over 5000 patients have received injectable granisetron hydrochloride in clinical trials.

Table 5 gives the comparative frequencies of the five commonly reported adverse events ($\geq 3\%$) in patients receiving granisetron hydrochloride injection, 40 mcg/kg, in single-day chemotherapy trials. These patients received chemotherapy, primarily cisplatin, and intravenous fluids during the 24-hour period following granisetron hydrochloride injection administration.

Table 5 Principal Adverse Events in Clinical Trials-Single-Day Chemotherapy

	Percent of Patients With Event	
	Granisetron HCl Injection* 40 mcg/kg (n=1268)	Comparator[†] (n=422)

Headache	14%	6%
Asthenia	5%	6%
Somnolence	4%	15%
Diarrhea	4%	6%
Constipation	3%	3%

*Adverse events were generally recorded over 7 days post-granisetron hydrochloride injection administration.

†Metoclopramide/dexamethasone and phenothiazines/dexamethasone.

In the absence of a placebo group, there is uncertainty as to how many of these events should be attributed to granisetron hydrochloride, except for headache, which was clearly more frequent than in comparison groups.

Radiation-Induced Nausea and Vomiting

In controlled clinical trials, the adverse events reported by patients receiving granisetron hydrochloride tablets and concurrent radiation were similar to those reported by patients receiving granisetron hydrochloride tablets prior to chemotherapy. The most frequently reported adverse events were diarrhea, asthenia, and constipation. Headache, however, was less prevalent in this patient population.

OVERDOSAGE

There is no specific treatment for granisetron hydrochloride overdosage. In case of overdosage, symptomatic treatment should be given. Overdosage of up to 38.5 mg of granisetron hydrochloride injection has been reported without symptoms or only the occurrence of a slight headache.

DOSAGE AND ADMINISTRATION

Emetogenic Chemotherapy

The recommended adult dosage of oral granisetron hydrochloride tablets is 2 mg once daily or 1 mg twice daily. In the 2 mg once-daily regimen, two 1 mg tablets are given up to 1 hour before chemotherapy. In the 1 mg twice-daily regimen, the first 1 mg tablet is given up to 1 hour before chemotherapy, and the second tablet 12 hours after the first. Either regimen is administered only on the day(s) chemotherapy is given. Continued treatment, while not on chemotherapy, has not been found to be useful.

Use in the Elderly, Pediatric Patients, Renal Failure Patients or Hepatically Impaired Patients : No dosage adjustment is recommended (see **CLINICAL PHARMACOLOGY: Pharmacokinetics**).

Radiation (Either Total Body Irradiation or Fractionated Abdominal Radiation)

The recommended adult dosage of oral granisetron hydrochloride tablets is 2 mg once daily. Two 1 mg tablets are taken within 1 hour of radiation.

Pediatric Use : There is no experience with oral granisetron hydrochloride tablets in the prevention of radiation-induced nausea and vomiting in pediatric patients.

Use in the Elderly : No dosage adjustment is recommended.

HOW SUPPLIED

Granisetron hydrochloride tablets 1 mg are white round film coated tablets debossed **cor** over **198** on one side and plain on the other side.

They are supplied as follows:

Blister cards of 2 unit of use tablets: NDC 64720-198-98

Blister cards of 20 unit of use tablets: NDC 64720-198-97 (intended for institutional use only)


Bottles of 20: NDC 64720-198-02 (intended for institutional use only)

Bottles of 100: NDC 64720-198-10 (intended for institutional use only)

Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].

Keep container closed tightly. Protect from light.

Keep this and all drugs out of the reach of children.

Manufactured and Distributed by:  **COREPHARMA**

Corepharma LLC

Middlesex, NJ 08846

MF # 489-02

August 2009

NDC 64720-198-02

COREPHARMA
Granisetron
HCl Tablets
1 mg*

Rx only

20 Tablets

***Each Tablet Contains:**
1.12 mg granisetron hydrochloride equivalent to
granisetron, 1 mg.

Usual Dosage: Chemotherapy: 2 mg once
daily. For patients with renal impairment,
regimen, two 1 mg tablets are given up to 1 hour
before chemotherapy. For patients with
renal impairment, the first 1 mg tablet is given up to 1 hour
before chemotherapy and the second tablet 12
hours after the first.

Radiotherapy: Two 1 mg tablets are taken within 1
hour of radiation. See accompanying prescribing
information.

Storage: Granisetron HCl (64720-198-02) is permitted to 15° to 30° C (59° to 86°) [see USP
Controlled Room Temperature].

**Granisetron HCl (64720-198-02) is not to be stored in the
USP. Keep container tightly closed, protect from
light.**

**KEEP THIS AND ALL DRUGS OUT OF THE
REACH OF CHILDREN.**

Manufactured and distributed by:
Corepharma, LLC
Madison, NJ 08446



MF # 627
Rev. 12/2007

Lot No.:
Exp. Date: